

PULL HERE TO OPEN ►

Paclo Pro[®]

ORNAMENTAL GROWTH REGULATOR

Active Ingredient:
Paclobutrazol
(±)-(R*,R*)-β-[(4-Chloro-phenyl)methyl]-α-(1,1-dimethylethyl)-1H-1,2,4-triazole-1-ethanol 00.4%

Other Ingredients: 99.6%

Total: 100.0%

Paclo Pro is formulated as an aqueous solution.

Paclo Pro contains 0.12 gram active ingredient per fluid ounce (4,000 ppm).

*EPA Reg. No. 100-996
EPA Est. No. 100-NE-001*

KEEP OUT OF REACH OF CHILDREN.

CAUTION

See additional precautionary statements and directions for use inside booklet.

*Product of United Kingdom
Formulated in the USA*

**SCP 996B-L1B 0814
4049640**

**1 gallon
Net Contents**

FIRST AID	
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment.	
HOT LINE NUMBER For 24-Hour Medical Emergency Assistance (Human or Animal) Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident), Call 1-800-888-8372	

PRECAUTIONARY STATEMENTS
<p>Hazards to Humans and Domestic Animals</p> <p style="text-align: center;">CAUTION</p> <p>Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Wear a long sleeved shirt and long pants, chemical resistant gloves made of any waterproof material (such as barrier laminate or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride or Viton®), and shoes plus socks. Remove and wash contaminated clothing before reuse.</p> <p>Personal Protective Equipment (PPE)</p> <p>Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category C on an EPA chemical resistance category selection chart.</p> <p>Applicators and other handlers must wear:</p> <ul style="list-style-type: none"> • Long-sleeved shirt and long pants • Chemical-resistant gloves made of any waterproof material, such as barrier laminate or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride or Viton® • Shoes plus socks <p>User Safety Requirements</p> <p>Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.</p>

continued...

PRECAUTIONARY STATEMENTS (*continued*)

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

For terrestrial uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

Physical and Chemical Hazards

Do not use or store near heat or open flame.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, LLC or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of this product contrary to label instructions or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and, (2) Buyer and User assume the risk of any such use. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.**

To the extent permitted by applicable law, in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Read all label directions carefully before use.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material, such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, polyvinyl chloride or viton.
- Shoes plus socks.

PRODUCT INFORMATION

Paclo Pro is a plant growth regulator for use on ornamental plants grown in containers in nurseries, greenhouses, shadehouses, and interiorscapes. Use of Paclo Pro effectively reduces internode elongation, resulting in more desirable compact plants. When used as directed, Paclo Pro produces no phytotoxic effects. **DO NOT REUSE POTS, TRAYS, OR OTHER CONTAINERS THAT PREVIOUSLY WERE USED IN THE PRODUCTION OF A CROP WHICH WAS TREATED WITH PACLO PRO.**

MIXING INSTRUCTIONS

Be sure the mix tank is clean and not contaminated with any material. Fill the spray tank with half the required amount of water. Using the dilution table (Table 1), determine the amount of Paclo Pro needed for the required concentration. Measure the desired volume accurately and add it to the tank.

Fill tank with the remaining amount of required water.

Agitate the mixture of Paclo Pro and water frequently to assure uniform distribution during application.

TABLE 1
Paclo Pro Dilution Table

ppm Paclo Pro Desired	Fl Oz Per Gallon	mL/Cc Per Gallon
0.05	0.0016	0.05
0.1	0.0032	0.1
0.2	0.0064	0.2
0.5	0.016	0.5
1	0.032	1.0
2	0.064	1.9
3	0.096	2.8
4	0.13	3.8
5	0.16	4.7
10	0.32	9.5
20	0.64	19.0
25	0.8	24.0
30	1.0	28.0
40	1.3	38.0
50	1.6	47.0
100	3.2	95.0
200	6.4	190.0

APPLICATION TECHNIQUES

Desired height control with Paclo Pro can be obtained with three different types of applications: sprays, drenches or bulb soaks.

Under certain conditions, sequential spray applications may be desirable.

Frequent agitation of the Paclo Pro solution and proper application techniques are critical in order to achieve desired results. Be sure of your calculations, volume measurements and sprayer calibration. When in doubt, recalculate.

1. Spray Applications

In spray applications, Paclo Pro penetrates into plant stems and is translocated to the terminal where it reduces internode elongation.

When applying a spray application, it is important that:

- Adequate spray volume is used to thoroughly wet plant stems. The misting technique used for some other growth regulators, where only upper leaves are covered with a light spray, will not produce desired results with Paclo Pro;
- Sprays are not applied to the point of excessive runoff into the potting media. The spray volume which drips down into the media may be desirable as it will be taken up by the roots and increase the effectiveness of Paclo Pro. However, too much runoff into the media may result in excessive height control;
- The spray technique provides thorough, consistent, uniform coverage of all plants. Failure to do so may result in non-uniform height control.

Paclo Pro may be applied at any time of the day without danger of burning leaves or causing chlorosis.

Overhead irrigation or rain 30 minutes after spray applications does not reduce the effectiveness of Paclo Pro.

Addition of wetting agents for spray applications is not necessary.

The recommended spray volume for small plants in small containers or plug trays which are closely spaced is 1-2 qt/100 sq ft of bench space. For larger plants with a well-developed canopy, a spray volume of 3 qt/100 sq ft of bench space is recommended.

Using up to 4 sequential applications at 2 week intervals may provide more uniform growth regulation and safety against over application. In general, sequential spray applications are to be applied using 50-100% of the lowest listed rate. Growers in cooler climates may have to use lower rates.

With some plant species, particularly chrysanthemums, hibiscus and azaleas, individual lateral shoots will outgrow the other laterals, causing non-uniform plant appearance. This results when individual laterals do not receive enough chemical when spray is applied. The use of sequential applications will reduce this problem.

2. Drench Applications

Application of Paclo Pro to the growing media will provide good control of plant height. Paclo Pro is readily absorbed by plant roots and translocated to the terminals.

Drench applications generally provide a longer-lasting, more uniform height control than spray applications, having little effect on flower size. Drench applications are very useful when applied late in the production cycle when plants have reached, or are near, the desired marketing size. Late drench applications are particularly useful on poinsettias, chrysanthemums and bulb crops.

Drench applications should be made to moist potting media. This may be achieved by watering plants the day before treatments. Drench applications to dry media will result in poor distribution.

Multiple plants growing in the same pot require a more uniform distribution of drench solution to achieve uniform height control.

Drench Rates and Volumes: The rates recommended for soil drench applications are based on a drench volume of 4 fl oz of final solution for an average 6-inch 'azalea' pot. Based on this recommendation, one gallon of solution will treat 32 6-inch pots. For smaller or larger pots, a suitable drench volume is enough final solution applied to achieve total run through of no more than 10%, providing that the potting media is properly moist before treatment. Table 2 may be used as a guide in determining appropriate drench volume needed for the specified pot sizes. For the grower who likes to apply Paclo Pro as a known amount of active ingredient per pot, Table 2 also shows the amount of active ingredient found in a specific volume at a known concentration.

TABLE 2
Guidelines and Conversions for Commonly Used Drench Volumes

Pot Diameter (inches)	Drench Volume (fl oz/pot)	Mg ai Paclo Pro/Pot			
		1 ppm	2 ppm	3 ppm	4 ppm
4"	2	0.063	0.125	0.188	0.25
5"	3	0.094	0.188	0.282	0.375
6"	4	0.125	0.25	0.375	0.50
8"	10	0.313	0.625	0.938	1.25
10"	25	0.783	1.56	2.35	3.125
10" hanging basket	15	0.470	0.939	1.41	1.878
12"	40	1.25	2.5	3.75	5.0

Note: The recommended drench volumes are based on the soil capacity of a common 6-inch 'azalea' type pot. Extrapolating the recommendation for this 6-inch 'azalea' type pot to smaller or larger containers may not be correct for total drench volume but should only be used as a guideline. The user must determine the appropriate rate and drench volume needed to achieve the desired result, based on both pot size and potting media used.

Paclo Pro can also be applied as a "drench" through sub-irrigation in saucers, benches or flooded floors. Using this method, the solution is applied to the media through the bottom of the container. Because most plant roots grow in the lower half of the container, this sub-application of Paclo Pro delivers the chemical to the plant more efficiently than the typical drench application, and therefore requires the use of lower rates than the typical drench. The optimum rates for a one-time sub-application is typically about 50-75% of the rate used in a typical drench. The optimum rate for continuous application in the irrigation water is about 10-33% of the rate needed for a one-time sub-irrigation application.

3. Preplant Bulb Soaks

Soaking of bulbs in solutions of Paclo Pro is also a very effective way to attain height control. The rates used and length of soaking time will vary, depending on the species. See the section on BULB CROPS for specific recommendations.

FACTORS AFFECTING PLANT RESPONSE TO PACLO PRO

IN ADDITION TO PROPER APPLICATION TECHNIQUE, THERE ARE SEVERAL ENVIRONMENTAL AND CULTURAL FACTORS WHICH CAN AFFECT A PLANT'S RESPONSE TO TREATMENT WITH PACLO PRO. These factors may cause a variation in the amount of Paclo Pro needed to provide desired plant height.

Cultural practices may affect the plant's response to Paclo Pro. Plants which are grown at close spacing or in smaller pots and using high water and fertilizer levels may require an increase in the amount of Paclo Pro needed.

For drench applications, plants grown in media with pine bark or a high organic content may require higher rates of Paclo Pro than those grown in media without pine bark or with a low organic content.

Different varieties or cultivars within a given plant species may require a higher or lower rate of Paclo Pro. The taller, more vigorous varieties generally require more chemical than do the naturally short, less vigorous varieties.

Growers should consult with plant and seed suppliers for vigor and other growth characteristics for newly released varieties.

Temperature can be the overriding factor in determining the amount of Paclo Pro needed. Stem elongation increases with increased temperatures. Growers in warm climates need to use higher rates and/or more applications compared to those in cooler climates.

The amount of Paclo Pro needed and number of applications may also vary depending on the time of year, with higher rates and/or more applications needed during warmer months.

DETERMINING OPTIMUM RATES

Optimum Paclo Pro rates will vary with different growers and will depend on their individual desired final plant height, growing conditions, and applications techniques. Different varieties or cultivars of the same species may respond differently to Paclo Pro. *Before Paclo Pro is applied to a large number of plants, growers should conduct trials with small numbers of plants using the recommended rates to determine the optimum rates for their situations.*

The rates recommended on this label are rate ranges and should be used only as guidelines. However, DO NOT exceed the maximum recommended rates.

The user should conduct trials on a small number of plants, adjusting the rate of Paclo Pro to achieve the desired height and length of control. For preplant bulk soak trials, it may be necessary to adjust both the rate and length of soak time in order to achieve desired results.

For plant species listed on the label, the user should run initial trials using the lowest recommended rates.

For plant species **not** specifically listed on the label, the user should run initial trials using the rates recommended in Table 3.

TABLE 3
Recommended Trial Rates (ppm) by General Plant Type*

Plant Type	Spray	Drench	Bulb Soak
Bedding Plants	30	1	N/A
Bedding Plant Plugs	5	NR	N/A
Flowering/Foliage Plants (Annual or Perennial)			
- Herbaceous Species	30	1	N/A
- Woody Species	50	2	N/A
Woody Landscape Plants	100	4	N/A
Bulb Crops	100	10	20 (@ 15 min)

NR = Use is not recommended

N/A = Use is not applicable

* The recommended trial rates are based on information developed primarily in the Sunbelt Region. Growers in regions north of the Sunbelt should run initial tests using half the recommended trial rates listed in Table 3.

USE AND RATE RECOMMENDATIONS BY CROP

Paclo Pro is effective in controlling the height of most ornamental crops. Be sure to read and fully understand the section on DETERMINING OPTIMUM RATES before applying to large numbers of plants.

A. Azaleas (Florist)

Paclo Pro can be used to control plant height, reduce bypass shoot elongation and promote flower bud initiation.

Spray applications are effective in the rate range of 100 to 200 ppm.

Drench Applications are effective in the rate range of 5 to 15 ppm. To control plant height and promote flower bud initiation, applications should begin when new growth, following final shaping, is 1-1/2 to 2 inches long.

To reduce bypass shoot development, applications should be made after bud set when bypass shoots are barely visible, or about 5 to 7 weeks prior to cooling.

B. Bedding Plants

Spray applications of Paclo Pro will provide height control of most bedding plants at a wide rate range of 5 to 90 ppm. The rate ranges for some specific bedding plants are:

Plant	Rate Range (ppm)	Plant	Rate Range (ppm)
Ageratum	15-45	Marigold (French)	15-30
Celosia	15-45	Pansy	5-15
Coleus	15-30	Petunia	15-45
Dahlia	15-45	Salvia	20-60
Dianthus	20-60	Snapdragon*	30-90
Impatiens (standard)	10-45	Verbena	15-30
Marigold (African)*	30-60	Zinnia	15-45

* Apply at an early stage of plant growth with good stem coverage, especially for vigorous varieties.

- Do not use on fibrous begonias as they are very sensitive to Paclo Pro. Overly stunted plants can result if they receive spray drift from applications made to surrounding species.
- Do not use on annual Vinca (periwinkle) as Paclo Pro may cause spotting of foliage, especially at high temperatures.

For bedding plants not specifically listed above, the user should determine optimum rates starting with a rate of 30 ppm in the Sunbelt Region and 15 ppm in the Northern Belt Region. Time of application should begin when new growth in height or width reaches 2 inches or when plants reach desired size to hold them at a marketable stage.

Late application timings and/or excessive rates may slow the growth of plants when transplanted. To avoid this, apply multiple applications at 1/4 to 1/2 the optimum rate.

High rates of Paclo Pro may delay flowering, especially of impatiens and petunias.

Drench applications are effective on bedding plants, but are recommended only for those plants in containers 6 inches or larger. The user should determine optimum rates, starting at 1 ppm.

C. Bedding Plant Plugs

Spray applications of Paclo Pro can also be used to effectively control height of bedding plant plugs. The recommended rate range of 1 to 20 ppm is much lower than the rate range for older bedding plants.

Plant	Rate Range (ppm)	Plant	Rate Range (ppm)
Ageratum	5-10	Marigold (French)	5-10
Celosia	5-10	Pansy	1-5
Coleus	5-10	Petunia	5-10
Dahlia	5-10	Salvia	5-10
Dianthus	10-20	Snapdragon	10-20
Impatiens (standard)	0.5-10	Verbena	5-10
Marigold (African)	10-20	Zinnia	5-10

For bedding plant plugs not specifically listed above, the user should determine optimum rates starting with a rate of 5 ppm. Timing of application should normally begin at the 1 to 2 true leaf stage.

Drench applications are **not** recommended for bedding plant plugs due to the sensitivity and extremely low rates needed.

For all uses of Paclo Pro on plugs, determining optimum rates should include an evaluation of the crop performance after transplanting to insure that treatment does not excessively reduce growth during the finished stage or in the landscape.

D. Bulb Crops

Height control can be achieved on a variety of bulb crops by any of the three application types.

Spray applications, although moderately effective, are the least desirable method for controlling height. Sequential applications are recommended in order to achieve desired uniformity. Applications should begin when plants are 2 to 4 inches tall.

Drench applications are very effective in the rate range of 8 to 160 ppm. Optimum rates vary widely, depending on species. Timing of application will also vary, depending on species. For bulbs which require a cold period, Paclo Pro is generally applied 1 to 5 days after removal from the cooler. For most other bulb types, application should be made when newly emerged shoots are 1 to 2 inches tall.

Preplant bulb soaks are also very effective. Effective rates for most species are in the range of 5 to 25 ppm, with a soaking time of 5-15 minutes. In general, lower use rates will require longer soaking times.

The following table gives recommended rate ranges and length of soaking time for a variety of bulb species:

Bulb Type	Spray Rate (ppm)	Drench Rate (ppm)	Preplant Bulb Soak Rate (ppm) / Soak Time
Amaryllis	ND	200	100 / 1 hr
Caladium	100-200	2-16	60 / 30 min
Calla Lily	ND	5-15	20 / 15 min
Dahlia	ND	10-40	>40 / 20 min
Freesia	ND	2-4	100-300 / 1 hr
Hybrid Lily	200-500	4-30	5-30 / 15 min
Montbretia	ND	ND	20-30 / 15 min
Tulip	ND	5-40	2-5 / 1 hr

> = Greater than.

ND = Rates for this particular use have not been determined. For these applications the user should run trials using the rates recommended in Table 3.

For species not specifically listed, trials should be conducted using rates outlined in the section on DETERMINING OPTIMUM RATES.

E. Chrysanthemums (Pot)

Paclo Pro is effective in controlling the height of pot chrysanthemums when applied as either a spray or drench.

Spray applications are effective at rates of 50 to 200 ppm. Applications should begin when axillary shoots are 2 to 3 inches long. Paclo Pro can be applied earlier to vigorous varieties if additional control is desired.

Sequential applications of lower rates generally provide more uniformly shaped plants than single-spray applications.

Drench applications of Paclo Pro are effective at recommended rates of 1 to 4 ppm.

Application timing during early production is when axillary shoots are 2 to 3 inches long.

Seasonably late applications are sometimes required at the time of disbud to prevent late stretch. Unlike spray applications, the drench can be safely applied with little to no effect on flowering.

Because pot chrysanthemums are usually planted with multiple cuttings per pot, uniform application is critical to achieving desired results.

F. Flowering Plants/Foliage Plants (not specifically listed)

Paclo Pro is effective as a spray or drench application in controlling height on a wide variety of other flowering plants and foliage plants. It can be used as either a holding agent to stop growth (e.g., interiorscape) or a toning agent to slow growth (i.e., when pot crops or hanging baskets are at or near marketable size). In general, herbaceous species will require lower rates than woody species. Trials should be conducted using rates outlined in the section on DETERMINING OPTIMUM RATES.

G. Geraniums

Geraniums are particularly sensitive to Paclo Pro. The user must determine optimum rates before applying to large numbers of plants.

Spray applications of Paclo Pro at recommended rates of 10 to 30 ppm will effectively control growth of geraniums. Early applications may require lower rates to avoid overstretching. Time of application for zonal geraniums is when growth is 1-1/2 to 2 inches long; for seed geraniums, 2 to 4 weeks after transplanting or when needed.

Paclo Pro will reduce late stretch when applied as the flower stems begin to elongate.

Drench applications, although effective, should be made with caution due to the extreme sensitivity of geraniums to Paclo Pro. Trials should be conducted to determine optimum rates.

H. Hibiscus

Spray applications at 30 to 150 ppm will effectively reduce shoot elongation. Application should be made when laterals are 1 to 4 inches long, depending on desired final plant size.

Single applications will control lateral growth for 3 to 6 weeks. Sequential applications may provide more uniform plant shape. Paclo Pro can be applied 1 to 2 weeks prior to flowering to prevent late stretch.

Drench applications will also effectively reduce shoot elongation. Trials should be conducted using recommended rates outlined in the section on DETERMINING OPTIMUM RATES.

I. Poinsettias

Spray applications of Paclo Pro will effectively control height of poinsettias. Recommended rates are 10 to 30 ppm for most areas of the U.S. In southern Florida, higher rates of 15 to 45 ppm are recommended.

Single applications may be made using the higher recommended rates. However, sequential applications initially using lower rates will provide better safety against overly retarded plants. For subsequent applications use 50 to 100% of the initial rate, depending on plant vigor at the time of reapplication.

Applications to slower growing varieties in cool climates should begin when axillary shoots are 2 to 3 inches long. For vigorous growing varieties in warm climates, applications should begin when axillary shoots are 1-1/2 to 2 inches long. Sequential applications may be applied 1 to 3 times, at 7 to 14-day intervals, depending on plant vigor/growth.

Seasonably late applications of Paclo Pro will reduce plant height, but, like most PGR's they may also reduce bract size. To prevent this, Paclo Pro should not be applied as a spray after initiation of short days. As a guide, do not apply Paclo Pro sprays after October 1 for areas outside of Florida, or after October 25 in Florida.

Drench applications are also an effective means of height control, and are most commonly used for late season application to plants which have initiated bracts or have reached or are near their desired marketing size.

Recommended rates are in the range of 0.25 to 3 ppm, based on a drench volume of 4 fl oz diluted solution per 6-inch pot. See MIXING INSTRUCTIONS section for directions on making the diluted solution.

Application during early production should be made when axillary shoots are 1-1/2 to 3 inches long. Early drenches of Paclo Pro may be made prior to or sequential to other PGR applications. As a guide, apply Paclo Pro at 0.05 ppm to 0.20 ppm. Vigorous Poinsettia varieties may require higher concentrations. When Paclo Pro is applied at low rates, bract formation and size are less affected. Subsequent growth may require additional drench applications every 10 to 21 days.

Seasonably late applications are sometimes required after initiation of short days to prevent late stretch and, unlike spray applications, can be safely applied with little to no effect on bract size.

NOTE: Optimum Paclo Pro rates and timings for both spray and drench applications to poinsettias will vary depending on the variety.

J. Woody Plants

Paclo Pro is effective in controlling height and initiating flower bud formation on a wide variety of woody plants using both spray or drench applications. Rate ranges for different species vary greatly. Trials should be conducted using rates outlined in the section on DETERMINING OPTIMUM RATES.

Examples of woody plants on which the product can be applied are:

Azalea	Hydrangea	Photinia
Bougainvillea	Ilex (Holly)	Pine
Camellia	Juniper	Rhododendron
Euonymus	Ligustrum	
Hibiscus	Magnolia	

USE DIRECTIONS FOR CHEMIGATION

In addition to the above use rates and recommendations, the following precautions must be observed when using this product in any type of irrigation system:

Apply this product only through the following systems:

- 1) Overhead sprinklers such as impact, micro-sprinklers, or booms.
- 2) Micro-irrigation such as spaghetti-tube or drip emitters.
- 3) Mist-type irrigation such as fog systems.
- 4) Hand-held calibrated equipment such as the hand-held wand with injector.
- 5) Sub-irrigation, such as ebb and flow and flooded floor systems, or through individual saucers.

Do not apply this product through any other type of irrigation system. Crop injury or lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water. If you have any questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Fill the supply tank with the desired amount of water. Then add the amount of Paclo Pro required in order to achieve the final solution rate recommended for the specific crop to be treated. Agitate the mixture of Paclo Pro and water frequently during the chemigation period to assure a uniform distribution throughout the system. Apply Paclo Pro continuously for the duration of the water application but do not exceed recommended rates and volumes as outlined on the product label. For overhead applications to the foliage and stems, apply at a volume of 1 to 2 qt per 100 sq ft for plugs and plants with small canopies. Volumes of 2 to 3 qt per 100 sq ft may be necessary for plants with large canopies. For applications to the soil, apply at a volume of 4 fl oz diluted solution per 6-inch pot. See MIXING INSTRUCTIONS section for directions on making the diluted solution.

SPRINKLER CHEMIGATION

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

CHEMIGATION SYSTEMS CONNECTED TO PUBLIC WATER SYSTEMS

Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days of the year.

Chemigation systems connected to public water systems must contain a functional, reduced pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water systems should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

The pesticide injection pipeline must contain a functional, automatic, quick closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where the pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage

Keep container closed when not in use. In case of spill or leak on floor or paved surfaces, soak up with sand, earth, or synthetic absorbent. Remove to chemical waste area.

Pesticide Disposal

Wastes resulting from the use of this product must be disposed of on-site or at an approved waste disposal facility.

Container Handling [less than or equal to 5 gallons]

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Paclo Pro® is a trademark of a Syngenta Group Company

Viton® is a trademark of E.I. duPont de Nemours and Company.

©2015 Syngenta

For non-emergency (e.g., current product information) call
Syngenta Crop Protection at 1-800-334-9481

Manufactured for:
Syngenta Crop Protection, LLC
P.O. Box 18300
Greensboro, North Carolina 27419-8300

**SCP 996B-L1B 0814
4049640**

Paclo Pro[®]

ORNAMENTAL GROWTH REGULATOR

Active Ingredient:	
Paclobutrazol (±)-(R*,R*)-β- [(4-Chlorophenyl)methyl]-α- (1,1-dimethylethyl)-1H-1,2,4- triazole-1-ethanol.	00.4%
Other Ingredients:	99.6%
Total:	100.0%

Paclo Pro is formulated as an aqueous solution.

Paclo Pro contains 0.12 gram active ingredient per fluid ounce (4,000 ppm). See additional precautionary statements and directions for use inside booklet.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

EPA Reg. No. 100-996
EPA Est. No. 100-NE-001
Paclo Pro[®] is a trademark of a Syngenta Group Company
©2015 Syngenta
Manufactured for:
Syngenta Crop Protection, LLC
P.O. Box 18300
Greensboro, North Carolina 27419-8300

**SCP 996B-L1B 0814
4049640**

1 gallon
Net Contents

KEEP OUT OF REACH OF CHILDREN. CAUTION

FIRST AID

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor or going for treatment.
HOT LINE NUMBER: For 24-Hour Medical Emergency Assistance (Human or Animal) Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident), Call **1-800-888-8372**.

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION

Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Wear a long sleeved shirt and long pants, chemical resistant gloves made of any waterproof material (such as barrier laminate or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride or Viton[®]), and shoes plus socks. Remove and wash contaminated clothing before reuse.

Environmental Hazards

For terrestrial uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

Physical and Chemical Hazards

Do not use or store near heat or open flame.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Keep container closed when not in use. In case of spill or leak on floor or paved surfaces, soak up with sand, earth, or synthetic absorbent. Remove to chemical waste area.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on-site or at an approved waste disposal facility.

Container Handling: Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

